

JOURNAL OF MATHEMATICAL ANALYSIS AND APPLICATIONS 35, 484-495 (1971)

Certain Properties of a Pair of Secondary Zeta-Functions

I. C. CHAKRAVARTY

Department of Mathematics, Trent University, Peterborough, Ontario, Canada

Submitted by Richard Bellman

1. INTRODUCTION

In a recent paper [2], we defined the "secondary zeta-functions" as a pair of functions derived from the Dirichlet series involving the powers of primes in one case and the non-trivial zeros of the Riemann zeta-function in the other and showed that the two secondary zeta-functions are related by a formula analogous to the functional equation of the Riemann zeta-function itself. This paper deals with a few properties of a pair of secondary zeta-functions.

For convenience of notation we make the convention throughout that $\frac{1}{2} + i\gamma$ shall run through the complex zeros of $\zeta(s)$, the Riemann zeta-function. Thus some values of γ will be complex if the Riemann hypothesis is false. Furthermore, if multiple zeros of $\zeta(s)$ exist, then the terms in sums over γ shall occur in their appropriate order of multiplicity.

The secondary zeta-functions $Z_p(s)$ and $Z_\gamma(s)$, whose properties we are going to investigate, are defined by:

$$Z_p(s) = \lim_{T \rightarrow \infty} \left\{ \sum_{0 < m \log p < T} \frac{\log p}{p^{\frac{1}{2}m}} (m \log p)^{-s} - \int_0^T e^{\frac{1}{2}u} u^{-s} du \right\},$$

where p runs through the prime numbers and m through the positive integers, and

$$Z_\gamma(s) = \sum_{\operatorname{Re}(\gamma) > 0} \gamma^{-s}.$$

The Dirichlet series involved in the definition of the function $Z_p(s)$ does not converge in the ordinary sense, but we can overcome this difficulty by using the notion of modified Abel summability which was used by Atkinson [1] to deal with certain divergent Dirichlet series in the theory of partitions.

The paper consists of four sections. In the section following this introduction, the poles of the function $Z_p(s)$ have been investigated. The behaviour of

the function $Z_\gamma(s)$ near $s = 1$ have been examined in the next section. Some particular values of the functions $Z_p(s)$ and $Z_\gamma(s)$ have been calculated in the last section.

2. THE POLES OF THE FUNCTION $Z_p(s)$

We have the Equation ([3], p. 58):

$$\frac{\zeta'}{\zeta}(s) = b - \frac{1}{s-1} - \frac{1}{2} \frac{\Gamma'}{\Gamma} \left(\frac{1}{2}s + 1 \right) + \sum_p \left(\frac{1}{s-\rho} + \frac{1}{\rho} \right), \quad (2.1)$$

where $b = \log 2\pi - \frac{1}{2}C - 1$, C being the Euler's constant, and $\rho = \frac{1}{2} + i\gamma$ runs through the nontrivial zeros of the function $\zeta(s)$.

Putting $s = \frac{1}{2} + t$ in (2.1), we obtain:

$$\frac{\zeta'}{\zeta} \left(\frac{1}{2} + t \right) = b - \frac{1}{t - \frac{1}{2}} - \frac{1}{2} \frac{\Gamma'}{\Gamma} \left(\frac{5}{4} + \frac{1}{2}t \right) + \sum_{\text{Re}(\gamma) > 0} \left(\frac{2t}{t^2 + \gamma^2} + \frac{1}{\frac{1}{4} + \gamma^2} \right). \quad (2.2)$$

We also have the result ([7], p. 247):

$$\frac{\Gamma'}{\Gamma}(1+z) = -C + \sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n+z} \right). \quad (2.3)$$

Writing $\frac{1}{4} + \frac{1}{2}t$ for z in (2.3), we obtain:

$$\frac{\Gamma'}{\Gamma} \left(\frac{5}{4} + \frac{1}{2}t \right) = -C + \sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n + \frac{1}{4} + \frac{1}{2}t} \right). \quad (2.4)$$

Equations (2.2) and (2.4) now give:

$$\begin{aligned} -\frac{\zeta'}{\zeta} \left(\frac{1}{2} + t \right) - \frac{1}{t - \frac{1}{2}} &= -b - \frac{1}{2}C + \frac{1}{2} \sum_{n=1}^{\infty} \left(\frac{1}{n} - \frac{1}{n + \frac{1}{4} + \frac{1}{2}t} \right) \\ &\quad - 2t \sum_{\text{Re}(\gamma) > 0} \frac{1}{t^2 + \gamma^2} - \sum_{\text{Re}(\gamma) > 0} \frac{1}{\frac{1}{4} + \gamma^2}. \end{aligned}$$

Therefore, if $f(t)$ represents the derivative with respect to t of the function on the left of the equation above, then

$$f(t) = \sum_{n=1}^{\infty} \frac{1}{(t + a)^2} + 2 \sum_{\text{Re}(\gamma) > 0} \frac{t^2 - \gamma^2}{(t^2 + \gamma^2)^2}, \quad (2.5)$$

where

$$a = 2n + \frac{1}{2}.$$

The function $f(t)$, thus defined, is regular at $t = \frac{1}{2}$ and consequently regular for all $t \geq 0$. As $t \rightarrow \infty$, it is of the order $O(t^{-2})$ and therefore, the function $F(s)$, the Mellin transform of $f(t)$, given by

$$F(s) = \int_0^\infty f(t) t^{s-1} dt \quad (2.6)$$

converges for $0 < \operatorname{Re}(s) < 2$. It is, however, possible to get an analytic continuation for $F(s)$.

Now, for $0 < \operatorname{Re}(s) < 2$, we have:

$$\int_0^\infty \frac{t^{s-1}}{(t+a)^2} dt = \frac{\pi(1-s) a^{s-2}}{\sin s\pi}, \quad (2.7)$$

and

$$\int_0^\infty \frac{t^2 - \gamma^2}{(t^2 + \gamma^2)^2} t^{s-1} dt = \frac{\pi(s-1) \gamma^{s-2}}{2 \sin \frac{1}{2}s\pi}. \quad (2.8)$$

Therefore, using (2.5) in (2.6), we have, with the help of (2.7) and (2.8):

$$\begin{aligned} F(s) &= \sum_{n=1}^\infty \frac{\pi(1-s) a^{s-2}}{\sin s\pi} + \sum_{\operatorname{Re}(\gamma) > 0} \frac{\pi(s-1) \gamma^{s-2}}{\sin \frac{1}{2}s\pi} \\ &= \frac{\pi(1-s)}{\sin s\pi} \chi(2-s) - \frac{\pi(1-s)}{\sin \frac{1}{2}s\pi} Z_\gamma(2-s), \quad (\operatorname{Re}(s) < 1), \end{aligned} \quad (2.9)$$

where

$$\chi(s) = 2^s \sum_{n=1}^\infty (4n+1)^{-s}, \quad (\operatorname{Re}(s) > 1) \quad (2.10)$$

and

$$Z_\gamma(s) = \sum_{\operatorname{Re}(\gamma) > 0} \gamma^{-s}, \quad (\operatorname{Re}(s) > 1).$$

Now, the functional equation connecting $Z_p(s)$ and $Z_\gamma(s)$ is given by [2]:

$$\begin{aligned} Z_p(s) - \Gamma(1-s) \{2^{1-s} - 2^{-s} \eta(1-s) - (2^{-s} - \frac{1}{2}) \zeta(1-s)\} \\ = -2\Gamma(1-s) \sin \frac{1}{2}s\pi Z_\gamma(1-s), \end{aligned} \quad (2.11)$$

where

$$\eta(s) = \sum_{n=0}^\infty (-1)^n (2n+1)^{-s}, \quad (\operatorname{Re}(s) > 1). \quad (2.12)$$

We may write

$$\eta(s) = \sum_{n=0}^{\infty} (4n+1)^{-s} - \sum_{n=0}^{\infty} (4n+3)^{-s},$$

so that

$$\begin{aligned} \sum_{n=1}^{\infty} (4n+1)^{-s} &= \sum_{n=0}^{\infty} (4n+1)^{-s} - 1 \\ &= \eta(s) + \sum_{n=0}^{\infty} (4n+3)^{-s} - 1 \\ &= \frac{1}{2}\eta(s) + \frac{1}{2} \sum_{n=0}^{\infty} (2n+1)^{-s} - 1 \\ &= \frac{1}{2}\eta(s) + \frac{1}{2}(1-2^{-s})\zeta(s) - 1, \end{aligned}$$

that is,

$$\frac{1}{2}\eta(s) + \frac{1}{2}(1-2^{-s})\zeta(s) = 1 + 2^{-s}\chi(s), \quad (\operatorname{Re}(s) > 1). \quad (2.13)$$

Replacing s by $1-s$, this can be written as:

$$2^{-s}\eta(1-s) + (2^{-s} - \frac{1}{2})\zeta(1-s) = 2^{1-s} + \chi(1-s), \quad (\operatorname{Re}(s) < 0). \quad (2.14)$$

Using (2.14), we can now write the functional Eq. (2.11) as:

$$Z_p(s) = -\Gamma(1-s)\chi(1-s) - 2\Gamma(1-s)\sin \frac{1}{2}s\pi Z_p(2-s), \quad (\operatorname{Re}(s) < 0). \quad (2.15)$$

Hence, with the help of (2.9), we can write:

$$\begin{aligned} Z_p(s-1) &= -\frac{\Gamma(2-s)\sin s\pi}{\pi(1-s)}F(s), \quad (\operatorname{Re}(s) < 1) \\ &= -\frac{F(s)}{\Gamma(s)} \\ &= -\frac{1}{\Gamma(s)} \int_0^{\infty} f(t) t^{s-1} dt, \quad (1 < \operatorname{Re}(s) < 2), \quad (2.16) \end{aligned}$$

using the analytic continuation of $F(s)$.

But, by definition,

$$f(t) = -\frac{d}{dt} \left\{ \frac{\zeta'}{\zeta} \left(\frac{1}{2} + t \right) \right\} + \frac{1}{(t - \frac{1}{2})^2}$$

Writing this for $f(t)$ and integrating by parts, (2.16) becomes:

$$= \frac{1}{\Gamma(s-1)} \int_0^{\infty} \left\{ \frac{\zeta'}{\zeta} \left(\frac{1}{2} + t \right) + \frac{1}{t - \frac{1}{2}} \right\} t^{s-2} dt,$$

since the integrated term vanishes for $1 < \operatorname{Re}(s) < 2$.

Hence,

$$Z_p(s) = \frac{1}{\Gamma(s)} \int_0^\infty \left\{ -\frac{\zeta'}{\zeta} \left(\frac{1}{2} + t \right) - \frac{1}{t - \frac{1}{2}} \right\} t^{s-1} dt, \quad (0 < \operatorname{Re}(s) < 1).$$

To get the analytic continuation for $Z_p(s)$, we write:

$$\begin{aligned} Z_p(s) &= \frac{1}{\Gamma(s)} \int_0^\infty \left\{ -\frac{\zeta'}{\zeta} \left(\frac{1}{2} + t \right) - \frac{1}{t - \frac{1}{2}} + \frac{1 - e^{1/2-t}}{t - \frac{1}{2}} \right\} t^{s-1} dt \\ &\quad + \frac{1}{\Gamma(s)} \int_0^\infty \frac{1 - e^{1/2-t}}{\frac{1}{2} - t} t^{s-1} dt \\ &= \frac{1}{\Gamma(s)} \int_0^\infty \left\{ -\frac{\zeta'}{\zeta} \left(\frac{1}{2} + t \right) - \frac{e^{1/2-t}}{t - \frac{1}{2}} \right\} t^{s-1} dt - \int_0^1 e^{\frac{1}{2}u} u^{-s} du, \end{aligned}$$

since

$$\begin{aligned} \int_0^\infty \frac{1 - e^{1/2-t}}{\frac{1}{2} - t} t^{s-1} dt &= - \int_0^\infty t^{s-1} dt \int_0^1 e^{(1/2-t)u} du \\ &= -\Gamma(s) \int_0^1 e^{\frac{1}{2}u} u^{-s} du, \quad (0 < \operatorname{Re}(s) < 1). \end{aligned}$$

We can now say that

$$Z_p(s) + \int_0^1 e^{\frac{1}{2}u} u^{-s} du$$

has a single valued analytic continuation for $\operatorname{Re}(s) > 0$ with no poles.

For $\operatorname{Re}(s) < 1$, we have

$$\int_0^1 e^{\frac{1}{2}u} u^{-s} du = \sum_{n=0}^{\infty} \frac{(\frac{1}{2})^n}{n!} \int_0^1 u^{n-s} du = \sum_{n=0}^{\infty} \frac{1}{2^n n! (n+1-s)}.$$

This shows that $\int_0^1 e^{\frac{1}{2}u} u^{-s} du$ has a continuation with simple poles of residues

$$-\frac{1}{2^{m-1}(m-1)!} \quad \text{at } s = m, \quad (m = 1, 2, 3, \dots).$$

Therefore, the function $Z_p(s)$ has simple poles of residues

$$\frac{1}{2^{m-1}(m-1)!} \quad \text{at } s = m, \quad (m = 1, 2, 3, \dots).$$

3. THE BEHAVIOUR OF THE FUNCTION $Z_p(s)$ NEAR $s = 1$

To begin with we assume the Riemann hypothesis to be correct and let $N(x)$ denote the number of zeros of the function $\zeta(s)$ in $0 < \operatorname{Im}(s) \leq x$.

Consider the function

$$T(s) = \int_0^{\infty} \left\{ N(x) - \left(\frac{x}{2\pi} \log \frac{x}{2\pi} - \frac{x}{2\pi} \right) \right\} x^{s-2} dx, \quad (3.1)$$

It is known ([6], p. 181) that the function

$$S(x) = N(x) - \left(\frac{x}{2\pi} \log \frac{x}{2\pi} - \frac{x}{2\pi} \right)$$

is of order $O(\log x)$ as $x \rightarrow \infty$. Hence, the integral (3.1) exists when x is infinite if $\operatorname{Re}(s) < 1$. As $x \rightarrow 0$, the integrand is of order $O(x^{s-1} \log x)$ and hence the integral exists at $x = 0$ if $\operatorname{Re}(s) > 0$. Therefore, the integral (3.1) converges for $0 < \operatorname{Re}(s) < 1$.

We write:

$$\begin{aligned} T(s) &= \int_1^{\infty} \left\{ N(x) - \left(\frac{x}{2\pi} \log \frac{x}{2\pi} - \frac{x}{2\pi} \right) \right\} x^{s-2} dx \\ &\quad - \frac{1}{2\pi} \int_0^1 \left(x^{s-1} \log \frac{x}{2\pi} - x^{s-1} \right) dx. \end{aligned}$$

The first integral on the right converges for $\operatorname{Re}(s) < 1$ and for $\operatorname{Re}(s) > 0$, the second integral is equal to

$$-\frac{1}{2\pi} \left\{ \frac{1 + \log 2\pi}{s} + \frac{1}{s^2} \right\}.$$

Therefore, we have:

$$\begin{aligned} T(s) &= \int_1^{\infty} \left\{ N(x) - \left(\frac{x}{2\pi} \log \frac{x}{2\pi} - \frac{x}{2\pi} \right) \right\} x^{s-2} dx \\ &\quad + \frac{1}{2\pi s^2} + \frac{1 + \log 2\pi}{2\pi s}, \quad (\operatorname{Re}(s) < 1). \end{aligned} \quad (3.2)$$

This will give us the analytic continuation of $T(s)$ for $\operatorname{Re}(s) < 0$. We can write:

$$\begin{aligned} T(s) &= \int_1^{\infty} N(x) x^{s-2} dx - \frac{1}{2\pi} \int_1^{\infty} \left(x \log \frac{x}{2\pi} - x \right) x^{s-2} dx \\ &\quad + \frac{1}{2\pi s^2} + \frac{1 + \log 2\pi}{2\pi s}, \quad (\operatorname{Re}(s) < 0). \end{aligned} \quad (3.3)$$

Integrating by parts, the second integral in (3.3) can be shown to be equal to

$$-\frac{1}{2\pi} \left\{ \frac{1}{s^2} + \frac{1 + \log 2\pi}{s} \right\}.$$

Therefore, Eq. (3.3) becomes:

$$\begin{aligned} T(s) &= \int_1^\infty N(x) x^{s-2} dx, \quad (\operatorname{Re}(s) < 0) \\ &= \frac{1}{1-s} \int_1^\infty x^{s-1} dN(x). \end{aligned}$$

Now, without the Riemann hypothesis, we have:

$$N(x) = \sum_{0 < \operatorname{Re}(\gamma) \leq x} 1$$

and so

$$T(s) = \frac{1}{1-s} \sum_{\gamma} \frac{1}{\{\operatorname{Re}(\gamma)\}^{1-s}} \quad (3.4)$$

Returning now to Equation (3.2), we obtain, using (3.4):

$$\frac{1}{1-s} \sum_{\gamma} \frac{1}{\{\operatorname{Re}(\gamma)\}^{1-s}} = \frac{1}{2\pi s^2} + \frac{1 + \log 2\pi}{2\pi s} + \int_1^\infty S(x) x^{s-2} dx, \quad (\operatorname{Re}(s) < 1).$$

Replacing s by $1-s$, we have:

$$\frac{1}{s} \sum_{\gamma} \frac{1}{\{\operatorname{Re}(\gamma)\}^s} = \frac{1}{2\pi(1-s)^2} + \frac{1 + \log 2\pi}{2\pi(1-s)} + \int_1^\infty S(x) x^{-s-1} dx, \quad (\operatorname{Re}(s) > 0).$$

As $x \rightarrow \infty$, $S(x) = O(\log x)$ and so the last integral is regular at $s = 1$. We can therefore write, near $s = 1$:

$$\sum_{\gamma} \frac{1}{\{\operatorname{Re}(\gamma)\}^s} = \frac{1}{2\pi(s-1)^2} - \frac{\log 2\pi}{2\pi(s-1)} + O(1) \quad (3.5)$$

But since $|\operatorname{Im}(\gamma)| < \frac{1}{2}$, we have:

$$\left| \frac{1}{\gamma^s} - \frac{1}{\{\operatorname{Re}(\gamma)\}^s} \right| = O\left(\frac{1}{\gamma^{s+1}}\right)$$

and so

$$\sum_{\gamma} \left\{ \frac{1}{\gamma^s} - \frac{1}{\{\operatorname{Re}(\gamma)\}^s} \right\}$$

would converge at $s = 1$ (as in $\sum_{\gamma} 1/\gamma^3$). This means that the difference is regular at $s = 1$. Hence, $Z_{\gamma}(s)$ behaves as the expression on the right-hand side of (3.5) near $s = 1$ and we may write:

$$Z_{\gamma}(s) = \frac{1}{2\pi(s-1)^2} - \frac{\log 2\pi}{2\pi(s-1)} + O(1). \quad (3.6)$$

We can now conclude that the function $Z_\gamma(s)$ has a double pole at $s = 1$ with residue $-(\log 2\pi)/2\pi$ and principal part

$$\frac{1}{2\pi(s-1)^2} - \frac{\log 2\pi}{2\pi(s-1)}.$$

4. SOME PARTICULAR VALUES OF THE FUNCTIONS $Z_p(s)$ AND $Z_\gamma(s)$

From Section 2 of this paper we have the functional equations [Eqs. (2.15) and (2.13)]:

$$Z_p(s) = -\Gamma(1-s)\chi(1-s) - 2\Gamma(1-s)\sin \frac{1}{2}s\pi Z_\gamma(1-s) \quad (4.1)$$

and

$$\chi(s) = 2^{s-1}\eta(s) + (2^{s-1} - \frac{1}{2})\zeta(s) - 2^s, \quad (4.2)$$

where $\chi(s)$ and $\eta(s)$ are given by (2.10) and (2.12) respectively.

The function $\eta(s)$ also satisfies a functional equation which can be derived from the general summation formula [5, p. 66]:

$$\sqrt{\alpha}\{f(\alpha) - f(3\alpha) + \cdots\} = \sqrt{\beta}\{F_s(\beta) - F_s(3\beta) + \cdots\}, \quad (4.3)$$

where $F_s(x)$ is the Fourier sine transform of $f(x)$ and $\alpha\beta = \pi/2$. If we let

$$f(x) = x^{-s} \quad (0 < \operatorname{Re}(s) < 1)$$

then

$$\begin{aligned} F_s(x) &= \sqrt{\frac{2}{\pi}} \int_0^\infty t^{-s} \sin xt \, dt \\ &= \sqrt{\frac{2}{\pi}} x^{s-1} \operatorname{Im} \left(\int_0^\infty e^{i u} u^{-s} \, du \right) \\ &= \sqrt{\frac{2}{\pi}} x^{s-1} \Gamma(1-s) \cos \frac{1}{2}s\pi. \end{aligned}$$

With these $f(x)$ and $F_s(x)$, the conditions of the summation formula are satisfied and the rightside of (4.3) becomes:

$$\sqrt{2\beta/\pi} \Gamma(1-s) \cos \frac{1}{2}s\pi \{\beta^{s-1} - (3\beta)^{s-1} + \cdots\},$$

Hence, using $\alpha\beta = \pi/2$, it follows from (4.3) that the functional equation for $\eta(s)$ can be written as:

$$\eta(s) = (\pi/2)^{s-1} \cos \frac{1}{2}s\pi \Gamma(1-s) \eta(1-s). \quad (4.4)$$

This gives:

$$\eta(0) = \frac{1}{2},$$

since

$$\eta(1) = 1 - \frac{1}{3} + \frac{1}{5} - \cdots = \frac{\pi}{4};$$

$$\eta(1 - 2m) = 0 \quad \text{and} \quad \eta(-2m) = \frac{1}{2} E_{2m},$$

since [4, p. 240]

$$\eta(2m + 1) = (-1)^m \frac{E_{2m}}{2^{2m+2}(2m)!} \pi^{2m+1}, \quad (4.5)$$

where E_{2m} is an Eulerian number.

We also have the following results [6, p. 19]:

$$\zeta(0) = -\frac{1}{2}, \quad \zeta(-2m) = 0$$

and

$$\zeta(1 - 2m) = \frac{(-1)^m B_{2m}}{2m},$$

where B_{2m} is a Bernoulli number.

Hence, we obtain from (4.2):

$$\chi(0) = -\frac{3}{4},$$

$$\chi(1 - 2m) = (-1)^m (2^{-2m} - \frac{1}{2}) \frac{B_{2m}}{2m} - 2^{1-2m}$$

and

$$\chi(-2m) = 2^{-2m} (\frac{1}{4} E_{2m} - 1).$$

We also know from Section 2 of this paper that the function $Z_p(s)$ has simple poles at $s = m$, ($m = 1, 2, 3, \dots$) with residues

$$\frac{1}{2^{m-1}(m-1)!}, \quad (m = 1, 2, 3, \dots).$$

Since $Z_p(s)$ has a simple pole at $s = 1$ with residue 1, we must have

$$\lim_{s \rightarrow 1} (s - 1) Z_p(s) = 1.$$

Hence, multiplying the functional Equation (4.1) throughout by $s - 1$ and taking limits of both sides $s \rightarrow 1$, we obtain:

$$\begin{aligned} 1 &= \{\chi(0) + 2Z_v(0)\} \lim_{s \rightarrow 1} \{1 - s\} \Gamma(1 - s) \\ &= \chi(0) + 2Z_v(0), \end{aligned}$$

whence

$$Z_v(0) = \frac{7}{8}.$$

In the same way we can calculate $Z_\nu(-2m)$. Considering the residue of $Z_\nu(s)$ at $s = 2m + 1$, we get:

$$\lim_{s \rightarrow (2m+1)} (s - 2m - 1) Z_\nu(s) = \frac{1}{2^{2m}(2m)!}.$$

Therefore, multiplying the functional Eq. (4.1) throughout by $s - 2m - 1$ and taking limits of both sides as $s \rightarrow 2m + 1$, we obtain:

$$\frac{1}{2^{2m}(2m)!} = \{\chi(-2m) + 2(-1)^m Z_\nu(-2m)\} \lim_{s \rightarrow 2m+1} \{(2m + 1 - s) \Gamma(1 - s)\}. \quad (4.6)$$

But

$$\begin{aligned} \lim_{s \rightarrow 2m+1} \{(2m + 1 - s) \Gamma(1 - s)\} &= \lim_{s \rightarrow 2m+1} \frac{(2m + 1 - s)\pi}{\Gamma(s) \sin s\pi} \\ &= \frac{1}{\Gamma(2m + 1)} = \frac{1}{(2m)!}, \end{aligned}$$

and so Eq. (4.6) gives:

$$Z_\nu(-2m) = (-1)^m \frac{8 - E_{2m}}{2^{2m+3}}.$$

In particular, using $E_2 = -1$, we get:

$$Z_\nu(-2) = -\frac{9}{32}.$$

To find $Z_\nu(0)$, we again consider the functional Eq. (4.1) and take limits of both sides as $s \rightarrow 0$. We obtain:

$$Z_\nu(0) = -\lim_{s \rightarrow 0} \{\chi(1 - s) + 2 \sin \frac{1}{2}s\pi Z_\nu(1 - s)\}, \quad (4.7)$$

where

$$\chi(1 - s) = 2^{-s}\eta(1 - s) + (2^{-s} - \frac{1}{2}) \zeta(1 - s) - 2^{1-s}.$$

Now, $\eta(1) = \pi/4$ and near $s = 0$, the function $\zeta(1 - s)$ can be written as ([6], p. 16):

$$\zeta(1 - s) = -\frac{1}{s} + C + O(s),$$

and

$$2^{-s} = 1 - s \log 2 + O(s^2).$$

Also, from the previous section we know that near $s = 0$, the function $Z_\nu(1 - s)$ can be written as (Eq. 3.6):

$$Z_\nu(1 - s) = \frac{1}{2\pi s^2} + \frac{\log 2\pi}{2\pi s} + O(1).$$

Hence, using the usual series expansion for $\sin \frac{1}{2} s\pi$, we can write (4.7) as:

$$\begin{aligned} 2 - \frac{\pi}{4} - \lim_{s \rightarrow 0} \left[\left\{ \frac{1}{2} - s \log 2 + O(s^2) \right\} \left\{ -\frac{1}{s} + C + O(s) \right\} \right. \\ \left. + \{s\pi + O(s^3)\} \left\{ \frac{1}{2\pi s^2} + \frac{\log 2\pi}{2\pi s} + O(1) \right\} \right] \\ = 2 - \left(\frac{\pi}{4} + \frac{1}{2} C + \frac{1}{2} \log 8\pi \right). \end{aligned}$$

But if we put $s = \frac{1}{2}$ in the known functional equation:

$$\frac{\zeta'}{\zeta}(s) = \log 2\pi + \frac{1}{2} \pi \cot \frac{1}{2} s\pi - \frac{\Gamma'}{\Gamma}(1-s) - \frac{\zeta'}{\zeta}(1-s),$$

we find

$$\frac{\zeta'}{\zeta}\left(\frac{1}{2}\right) = \frac{\pi}{4} + \frac{1}{2} C + \frac{1}{2} \log 8\pi.$$

Therefore

$$Z_p(0) = 2 - \frac{\zeta'}{\zeta}\left(\frac{1}{2}\right).$$

To find $Z_p(-2m)$, we observe that the function $Z_p(s)$ is regular at $s = 2m + 1$, ($m = 1, 2, 3, \dots$) and therefore the functional Eq. (4.1) should give us values of $Z_p(-2m)$, ($m = 1, 2, 3, \dots$). Putting $s = -2m$ in the afore-said functional equation, we obtain:

$$\begin{aligned} Z_p(-2m) &= -(2m)! \{2^{2m} \eta(2m+1) + (2^{2m} - \tfrac{1}{2}) \zeta(2m+1) - 2^{2m+1}\} \\ &= (2m)! \{2^{2m+1} - (2^m - \tfrac{1}{2}) \zeta(2m+1)\} - \tfrac{1}{4} (-1)^m \pi^{2m+1} E_{2m}, \end{aligned}$$

using (4.5).

In particular, using $\eta(3) = \pi^3/32$, we get:

$$Z_p(-2) = 16 - \frac{\pi^3}{4} - 7\zeta(3).$$

REFERENCES

1. F. V. ATKINSON, The abel summation of certain dirichlet series, *Quart. J. Math. Oxford Ser.* **19** (1948), 59-64.
2. I. C. CHAKRAVARTY, The secondary zeta-functions, *J. Math. Anal. Appl.* **30** (1970), 280-294.
3. A. E. INGHAM, "The Distribution of Prime Numbers," Cambridge Tracts in Mathematics and Mathematical Physics, No. 30, Cambridge University Press, London, 1932.

4. K. KNOPP, "Theory and Application of Infinite Series," 2nd Ed., Blackie and Son Ltd., London and Glasgow, 1951.
5. E. C. TITCHMARSH, "Introduction to the Theory of Fourier Integrals," 2nd Ed., Oxford, Clarendon Press, 1948.
6. E. C. TITCHMARSH, "The Theory of the Riemann Zeta-Function," Oxford, Clarendon Press, 1951.
7. E. T. WHITTAKER AND G. N. WATSON, "A Course of Modern Analysis," 4th Ed., Cambridge University Press, 1962.